

Letter to the Editor

Strategic Decision-Making in Elderly Healthcare Management: Integrating Cognitive, Motivational, and Environmental Factors

Maryam Moghimian¹, Fahimeh Kashani², Masoumeh Rostami³, Sasan Navkhasi^{3,2}

¹Department of Nursing, Na.C., Islamic Azad University, Najafabad, Iran

²Community health research center, Isf.C., Islamic Azad University, Isfahan, Iran

³Department of Nursing, Asadabad School of Medical Sciences, Asadabad, Iran

*Corresponding author: sasan Navkhasi, Email: sasan.navkhasi@gmail.com



Please cite this article as follows: Moghimian M, Kashani F, Rostami M, Navkhasi S, Navkhasi S. Strategic decision-making in elderly healthcare management: integrating cognitive, motivational, and environmental factors. Avicenna J Aging Healthc 2025;3(1):47-49. doi:10.34172/ajah.4034

Received: June 6, 2026, Accepted: August 8, 2026, ePublished: August 29, 2026

The accelerating growth of the global elderly population is a significant challenge for the 21st century (1). Contemporary societies are confronted with multifaceted and complex challenges stemming from the psychological, social, cultural, and economic dimensions of aging, imposing a significant burden on both families and nations (2). This life stage is marked by concomitant physical, physiological, motor, and cognitive transformations (3). Demographic trends in Iran illustrate a rapid demographic shift; in other words, census data show that the population aged 60 and above rose from approximately 2.8% in 2011 to about 26.9% in 2016, with projections estimating that it will reach 25% by 2051 (4). Consequently, a comprehensive investigation into the ramifications of population aging is imperative to equip policymakers with the insights necessary to strategically address both the opportunities and challenges associated with this transition (5). Addressing the specific issues and needs of this demographic is thus a critical societal imperative (6). In deconstructing this imperative, the subsequent analysis examines the cognitive, environmental, and motivational factors essential for strategic healthcare planning for the elderly.

Cognitive Factors

Cognitive decline in older adults manifests through symptoms including memory impairment, diminished capacity in performing activities of daily living, and behavioral alterations such as aggression or agitation. These deficits typically progress with age and exert substantial impacts on personal and social functioning (7). Management strategies for cognitive health should

incorporate regular physical activity, structured exercise regimens, and physical training. These interventions aid in preserving and enhancing cognitive functions such as memory and concentration by promoting cerebral blood circulation, fortifying neural connectivity, alleviating stress, and fostering neuroplasticity (8). Moreover, sustained social integration, achieved through interpersonal interactions, club memberships, participation in group-based leisure activities (e.g., chess, card games, etc.), prosocial behaviors, civic engagement, volunteerism, and lifelong learning, is correlated with a markedly reduced risk of cognitive deterioration. It is important to note that impaired mobility, often requiring the use of assistive devices, can present a significant barrier to such beneficial social participation (9, 10). Adherence to dietary patterns such as the Mediterranean diet, as endorsed by the World Health Organization, is also recognized as beneficial. This diet, characterized by high intake of fresh fruits, vegetables, legumes, nuts, whole grains, omega-3-rich fish, and olive oil, alongside the consumption of vibrantly colored fruits and vegetables like berries, leafy greens, and tomatoes, provides essential vitamins and antioxidants that exert significant neuroprotective effects (11).

Environmental Factors

Older adults represent a demographic group particularly vulnerable to the detrimental impacts of climate change. Exposure to extreme ambient temperatures is associated with elevated mortality rates, especially among those with pre-existing chronic conditions and lower socioeconomic status. Concurrently, age-related physiological decline, including a reduced capacity to metabolize environmental



toxins and diminished respiratory function, heightens susceptibility to air pollution. This vulnerability is characterized by increased rates of respiratory and cardiovascular pathologies, as well as an accelerated decline in cognitive performance. Empirical studies substantiate this relationship; for instance, a 1°C increase in annual mean temperature was linked to a 3% reduction in hypertension, a 6% reduction in cardiac disease, and a 5% reduction in stroke among individuals aged over 65 in China, alongside a general rise in emergency hospital admissions within this cohort (1). Among environmental pollutants, particulate matter (PM) demonstrates the most pronounced adverse effects on composite aging indices, being associated with accelerated phenotypic aging and reduced cerebral volume. Furthermore, exposure to noise pollution and limited access to green spaces have been correlated with decreased gray matter volume and exacerbated brain aging (12). Mitigating the environmental impact on the elderly necessitates a comprehensive multi-pronged strategy. This includes implementing robust systems for monitoring air, water, and soil quality, accelerating the transition to clean and renewable energy sources, and enhancing water and sanitation infrastructure. The safe management of chemicals and remediation of polluted sites are fundamental components of this approach. In the housing sector, quality improvements through thermal insulation and the installation of sustainable heating and cooling systems are crucial for maintaining thermal comfort. At the service delivery level, establishing early warning systems for extreme weather events and creating local support centers (e.g., cooling centers in summer, warming centers in winter) enable timely interventions. Complementary public awareness campaigns educating older adults and their caregivers on coping strategies for temperature extremes, including adequate hydration and appropriate clothing, are integral parts of this holistic action plan (13, 14).

Motivational Factors

A primary manifestation of diminished motivation among elderly individuals, particularly those requiring care and diagnosed with depression, includes a marked reduction in energy levels, apathy, listlessness, anhedonia, emotional blunting, social isolation, and neglect of daily routines. These symptoms are frequently compounded by physical health issues such as chronic pain, functional limitations, and the fear of becoming dependent, resulting in a significant decline in self-initiated and goal-oriented behavior (15). Therefore, understanding the motivational drivers, including health concerns and personal goals, that crucially shape societal expectations for active aging is of paramount importance (16). Key motivators can be categorized into 3 domains: medical (e.g., health maintenance, back pain prevention, improved joint mobility, etc.), participatory (e.g., desire for social connection, receiving social support, etc.), and physical-psychological (e.g., enhancing physical fitness, stress reduction, etc.) (6, 17). When effectively harnessed through interventions such as physical exercise,

these motivational factors can contribute to increased longevity, enhanced self-confidence and autonomy, strengthened social bonds, reduced feelings of loneliness, and ultimately, the alleviation of psychological conditions such as depression (2).

Strategic decision-making for elderly healthcare management necessitates the synergistic integration of cognitive, environmental, and motivational factors. This integrated paradigm paves the way for the formulation of evidence-based and responsive healthcare strategies by addressing the complexities of demographic aging and the multidimensional needs of the elderly population.

Authors' Contribution

Conceptualization: Maryam Moghimian, Sasan Navkhasi
Data Curation: Masoumeh Rostami
Formal Analysis: Maryam Moghimian, Fahimeh Kashani
Funding Acquisition: Sasan Navkhasi
Investigation: Maryam Moghimian
Methodology: Maryam Moghimian, Sasan Navkhasi
Project Administration: Maryam Moghimian, Sasan Navkhasi
Resources: Masoumeh Rostami, Sasan Navkhasi
Software: Masoumeh Rostami, Sasan Navkhasi
Supervision: Maryam Moghimian, Fahimeh Kashani
Validation: Maryam Moghimian, Fahimeh Kashani
Visualization: Maryam Moghimian, Fahimeh Kashani
Writing–Original Draft: Maryam Moghimian, Fahimeh Kashani
Writing–Review & Editing: Maryam Moghimian, Sasan Navkhasi

Competing Interests

None declared.

Ethical Approval

Not applicable.

Funding

None declared.

References

1. Zanjari N, Mollaei P. The effects of climate change on the health of the older adults: a systematic review. *Iran J Health Environ* 2024;17(2):397-416.
2. Ahmadi M, Lida F, Dovin F, Mirzazadeh, Zahrasadat, Peymani Zad. An Analytical Study of the Social and Motivational Effects of Elderly Sports in Iran. *Strategic Sociological Studies in Sport*. 2023;9(3):403-14.
3. Masdari V, Sheikh M, Hoomanian D, Ghayournajafabadi M, Bagherzadeh F. The effect of indoor rock climbing with and without environmental affordances on balance and fear of falling in older. *Sci J Rehabil Med* 2025;13(6):1108-21. doi:10.32598/SJRM.13.6.3273
4. Mansouri-Shad F, Cheraghi P, Mafakheri-Bashmagh S, Nouri Q, Cheraghi Z. Assessing the elderly health-related unmet needs and some related factors in Hamadan city, Iran. *Iran J Ageing* 2025;20(1):52-69. doi:10.32598/sija.2024.1202.3
5. Bahador F, Mahfoozpour S, Masoudi Asl I, Vahdat S. Identifying the main factors of providing primary preventive care services to the elderly in Iran. *Iran J Ageing* 2023;17(4):506-21. doi:10.32598/sija.2022.3180.3
6. Kholousi P, Tojari F, Esmaeili Shazandi MR. Determining motives and motivation factors among Iranian elders of male and female to participate in physical activities. *Strategic Studies on Youth and Sports* 2018;17(40):27-42.
7. Borji S, Rahaei Z, Aminisani N. Demographic determinants of the cognitive status among older adults: Neyshabur longitudinal study on aging. *The Journal of Tolooebehdasht* 2024;23(1):67-83. doi:10.18502/tbj.v23i1.15524

8. Saadatmehr SR, Akhavi pour A, Pourmoradkohan P, Bakhshian F, Amiri M. The effect of brain gym training on cognitive abilities of the elderly. *Journal of Sport Management and Motor Behavior* 2024;19(38):135-66. doi:10.22080/jsmb.2024.17232.3215
9. Yang Q, Lin S, Zhang Z, Du S, Zhou D. Relationship between social activities and cognitive impairment in Chinese older adults: the mediating effect of depressive symptoms. *Front Public Health* 2024;12:1506484. doi:10.3389/fpubh.2024.1506484
10. Borji M, Moradi M, Mahdian K, Bahmanpour K, Navkhasi S, Salimi E. Comparison of social participation level between older adults with assistive mobility devices and those without these devices. *Iran J Rehabil Res Nurs* 2016;3(2):38-44.
11. World Health Organization (WHO). Risk Reduction of Cognitive Decline and Dementia: WHO guidelines. WHO; 2019. Available from: <https://iris.who.int/handle/10665/312180>.
12. Pu F, Chen W, Li C, Fu J, Gao W, Ma C, et al. Heterogeneous associations of multiplexed environmental factors and multidimensional aging metrics. *Nat Commun* 2024;15(1):4921. doi:10.1038/s41467-024-49283-0
13. Luo H, Hu H, Zheng Z, Sun C, Yu K. The impact of living environmental factors on cognitive function and mild cognitive impairment: evidence from the Chinese elderly population. *BMC Public Health* 2024;24(1):2814. doi:10.1186/s12889-024-20197-2
14. Landrigan PJ, Fuller R, Acosta NJ, Adeyi O, Arnold R, Basu NN, et al. The Lancet Commission on pollution and health. *Lancet* 2018;391(10119):462-512. doi:10.1016/s0140-6736(17)32345-0
15. Vathke C, Kessler EM, Gellert P, O'Sullivan JL, Lech S. Motivation and psychotherapy in older adults with care needs and depression. *GeroPsych* 2023;36(2):49-62. doi:10.1024/1662-9647/a000305
16. Radoš S, Pavlova MK, Rothermund K, Silbereisen RK. Perceived expectations for active aging: the role of motivational and personality factors. *Aging Ment Health* 2024;28(11):1559-66. doi:10.1080/13607863.2024.2385450
17. Khoukhazade T, Navkhasi S, Khalafzade A, Moradi M, Veysi A, Borji M. The relationship between physical activity and balance control in the elderly with diabetes in Ilam, Iran, in 2016. *J Diabetes Nurs* 2016;4(3):60-72.